

APPROXIMATING REALS BY SUMS OF RATIONALS

TSZ HO CHAN AND ANGEL V. KUMCHEV

ABSTRACT. We study how well a real number can be approximated by sums of two or more rational numbers with denominators up to a certain size.

1. INTRODUCTION

Dirichlet's theorem on diophantine approximation tells us that we can approximate any real number by rational numbers quite well, namely:

Theorem 1. *For any real θ and any positive integer N , there exist integers a and q , with $1 \leq q \leq N$, such that*

$$\left| \theta - \frac{a}{q} \right| < \frac{1}{qN}.$$

Moreover, the bound $1/(qN)$ is best possible, apart from the constant factor. To see this, it suffices to consider the golden ratio $\theta = (\sqrt{5}-1)/2$ (see [3, §11.8]). During his work in [1], the first author accidentally stumbled across the following analogous question:

Question 1. *For any real θ and any positive integer N , give an upper bound for*

$$\min_{\substack{a_1, a_2, q_1, q_2 \in \mathbb{Z} \\ 1 \leq q_1, q_2 \leq N}} \left| \frac{a_1}{q_1} + \frac{a_2}{q_2} - \theta \right|.$$

With the golden ratio in mind, we know that the upper bound can be no better than $O(1/(q_1 q_2 N^2))$. So, what is the best possible upper bound? More generally,

Question 2. *Let k be a positive integer. For any real θ and any positive integer N , give an upper bound for*

$$\min_{\substack{a_1, \dots, a_k, q_1, \dots, q_k \in \mathbb{Z} \\ 1 \leq q_1, \dots, q_k \leq N}} \left| \frac{a_1}{q_1} + \dots + \frac{a_k}{q_k} - \theta \right|.$$

To these, we have the following result:

2000 *Mathematics Subject Classification.* 11J04, 11K60.

Theorem 2. *Let k be a positive integer. For any real θ and any positive integer N , there exist integers $a_1, \dots, a_k, q_1, \dots, q_k$, with $1 \leq q_1, \dots, q_k \leq N$, such that*

$$\left| \frac{a_1}{q_1} + \dots + \frac{a_k}{q_k} - \theta \right| \ll N^{-k}.$$

The bound N^{-k} is best possible in the sense that, for some θ , the minimum in Question 2 can be as large as N^{-k} . For example, if one considers $\theta = 1/(2N^k)$,

$$\left| \frac{a_1}{q_1} + \dots + \frac{a_k}{q_k} - \theta \right| \geq \frac{1}{2N^k}$$

for any choice of $a_1, \dots, a_k, q_1, \dots, q_k$, with $1 \leq q_1, \dots, q_k \leq N$. However, one expects such pathological examples to be relatively rare, and so one may wonder if it is possible to obtain a sharper upper bound involving the q_i 's. For example, is it possible to replace N^{-k} by $(q_1 \cdots q_k)^{-1} N^{-k}$ in Theorem 2? We shall briefly address this issue in the last section.

2. P T 2

Lemma. *Suppose that $k \geq 1$ is an integer. There is a number $x_0(k) \geq 1$ such that*

$$\sum_{1 \leq q_1, \dots, q_k \leq x}^* q_1 \cdots q_k \gg x^{2k},$$

whenever $x \geq x_0(k)$. Here, $\sum^*$ denotes a summation over the k -tuples $q_1, \dots, q_k$ such that $(q_i, q_j) = 1$ whenever $1 \leq i < j \leq k$.

Proof. It suffices to show that

$$\sum_{\substack{n \leq x \\ (n, m) = 1}} \phi(n)^\alpha n^{1-\alpha} \gg x^2 \phi(m) m^{-1}, \quad (1)$$

whenever $0 \leq \alpha \leq k-1$, $1 \leq m \leq x^{k-1}$, and $x \geq x_0(k)$. The conclusion of the lemma will then follow by successive applications of (1) with $\alpha = 0, 1, \dots, k-1$ to the summations over $q_k, q_{k-1}, \dots, q_1$.

We now proceed to establish (1). We start by showing that

$$\sum_{\substack{n \leq x \\ (n, m) = 1}} \left(\frac{n}{\phi(n)} \right)^\alpha \ll x \phi(m) m^{-1}. \quad (2)$$

Define the multiplicative functions

$$f(n) = \begin{cases} (n/\phi(n))^\alpha & \text{if } (n, m) = 1, \\ 0 & \text{if } (n, m) > 1, \end{cases} \quad g(n) = \sum_{d|n} f(d) \mu(n/d).$$

Then $g(n) \geq 0$, and

$$\begin{aligned}
\sum_{n \leq x} f(n) &= \sum_{n \leq x} \sum_{d|n} g(d) = \sum_{d \leq x} g(d) \left\lfloor \frac{x}{d} \right\rfloor \leq x \sum_{d \leq x} g(d) d^{-1} \\
&\leq x \prod_{p \leq x} \sum_{v=0}^{\infty} g(p^v) p^{-v} = x \prod_{p \leq x} (1 - p^{-1}) \sum_{v=0}^{\infty} f(p^v) p^{-v} \\
&\leq x \prod_{\substack{p|m \\ p \leq x}} (1 - p^{-1}) \prod_p \left(1 + \frac{p^\alpha - (p-1)^\alpha}{p(p-1)^\alpha} \right) \\
&\leq x \prod_{p|m} (1 - p^{-1}) \prod_p \left(1 + \frac{p^\alpha - (p-1)^\alpha}{p(p-1)^\alpha} \right) + O(1).
\end{aligned}$$

The last inequality follows on noting that m has at most $k-2$ prime divisors $p > x$, and hence,

$$\prod_{\substack{p|m \\ p > x}} (1 - p^{-1}) = 1 + O(x^{-1}).$$

This proves (2). On the other hand, when $\alpha = 0$, we have

$$\sum_{\substack{n \leq x \\ (n,m)=1}} n = \sum_{d|m} \mu(d) d \sum_{k \leq x/d} k = \frac{\phi(m)}{2m} x^2 + O(x\tau(m)),$$

whence

$$\sum_{\substack{n \leq x \\ (n,m)=1}} n^{1/2} \geq x^{-1/2} \sum_{\substack{n \leq x \\ (n,m)=1}} n \gg x^{3/2} \phi(m) m^{-1}. \quad (3)$$

Finally, (1) follows from (2), (3), and Cauchy's inequality:

$$\sum_{\substack{n \leq x \\ (n,m)=1}} \phi(n)^\alpha n^{1-\alpha} \geq \left\{ \sum_{\substack{n \leq x \\ (n,m)=1}} n^{1/2} \right\}^2 \left\{ \sum_{\substack{n \leq x \\ (n,m)=1}} \left(\frac{n}{\phi(n)} \right)^\alpha \right\}^{-1} \gg x^2 \phi(m) m^{-1}.$$

□

Proof of Theorem 2. For $0 < \Delta < 1/2$, define

$$t(x) = \max(1 - |x|/\Delta, 0), \quad g(x) = \sum_{n=-\infty}^{\infty} t(x-n).$$

The function g has a Fourier expansion

$$g(x) = \sum_{h=-\infty}^{\infty} \hat{g}_h e(hx), \quad \hat{g}_h = \Delta \left(\frac{\sin \pi \Delta h}{\pi \Delta h} \right)^2.$$

We consider the sum

$$\mathcal{S} = \sum_{1 \leq q_1, \dots, q_k \leq N}^* \sum_{a_1=1}^{q_1} \cdots \sum_{a_k=1}^{q_k} g \left(\frac{a_1}{q_1} + \cdots + \frac{a_k}{q_k} - \theta \right), \quad (4)$$

where $\sum^*$ has the same meaning as in the Lemma. Putting in the Fourier expansion for g , we get

$$\begin{aligned} \mathcal{S} &= \sum_{1 \leq q_1, \dots, q_k \leq N}^* \sum_{a_1=1}^{q_1} \cdots \sum_{a_k=1}^{q_k} \sum_{h=-\infty}^{\infty} \hat{g}_h e \left(h \left(\frac{a_1}{q_1} + \cdots + \frac{a_k}{q_k} - \theta \right) \right) \\ &= \sum_{1 \leq q_1, \dots, q_k \leq N}^* \sum_{h=-\infty}^{\infty} \hat{g}_h e(-h\theta) \sum_{a_1=1}^{q_1} e(ha_1/q_1) \cdots \sum_{a_k=1}^{q_k} e(ha_k/q_k) \\ &= \sum_{1 \leq q_1, \dots, q_k \leq N}^* q_1 \cdots q_k \sum_{\substack{h=-\infty \\ q_1 \cdots q_k | h}}^{\infty} \hat{g}_h e(-h\theta), \end{aligned} \quad (5)$$

as

$$\sum_{a=1}^q e(ha/q) = \begin{cases} q & \text{if } q \mid h, \\ 0 & \text{otherwise.} \end{cases}$$

If m is a positive integer and $\Delta \leq m^{-1}$, we have

$$\begin{aligned} \sum_{h \neq 0} |\hat{g}_{mh}| &\leq 2 \left\{ \sum_{h=1}^H \Delta + \frac{1}{\Delta m^2} \sum_{h=H+1}^{\infty} h^{-2} \right\} \\ &\leq 2 \left(H\Delta + \frac{1}{H\Delta m^2} \right) \leq 6m^{-1}, \end{aligned} \quad (6)$$

where $H = \lceil (\Delta m)^{-1} \rceil$; whereas if $\Delta > m^{-1}$, we have

$$\sum_{h \neq 0} |\hat{g}_{mh}| \leq \frac{2\zeta(2)}{\Delta m^2} \leq 4m^{-1}. \quad (7)$$

Putting (6) and (7) (with $m = q_1 \cdots q_k$) into (5), we obtain

$$\mathcal{S} = \Delta \sum_{1 \leq q_1, \dots, q_k \leq N}^* q_1 \cdots q_k + O(N^k),$$

the O -implied constant being absolute (in fact, it is 6). Therefore, upon choosing $\Delta = cN^{-k}$ with a sufficiently large $c > 0$, it follows from the Lemma that $\mathcal{S} > 0$. Hence, by (4),

$$g \left(\frac{a_1}{q_1} + \cdots + \frac{a_k}{q_k} - \theta \right) > 0$$

for some integers $a_1, \dots, a_k, q_1, \dots, q_k$ with $1 \leq q_1, \dots, q_k \leq N$. Then, by the definition of g ,

$$\left| \frac{a_1}{q_1} + \dots + \frac{a_k}{q_k} - n - \theta \right| \leq cN^{-k}$$

for some integer n . This establishes the theorem. $\square$

3. C

We conclude this note with a short discussion of possible improvement on the bound N^{-k} in Theorem 2. For example, is it possible to replace N^{-k} by $(q_1 \cdots q_k)^{-1} N^{-k}$? While such a result may appear to be the right generalization of Dirichlet's theorem, it is not true in general. Indeed, suppose that for any real θ , there exist integers $a_1, \dots, a_k, q_1, \dots, q_k$, with $1 \leq q_1, \dots, q_k \leq N$, such that

$$\left| \frac{a_1}{q_1} + \dots + \frac{a_k}{q_k} - \theta \right| \leq \frac{C}{q_1 \cdots q_k N^k}. \quad (8)$$

Then

$$[0, 1] \subseteq \bigcup_{q \in \mathcal{D}_k(N)} \bigcup_{0 \leq a \leq q} \{ \theta \in \mathbb{R} : |\theta - a/q| \leq C/(qN^k) \}, \quad (9)$$

where $\mathcal{D}_k(N)$ denotes the set of least common denominators of the sums appearing on the left side of (8). By a result of Erdős [2], $\mathcal{D}_k(N)$ has cardinality

$$|\mathcal{D}_k(N)| \ll N^k (\log N)^{-c}$$

for some constant $c = c(k) > 0$, so it follows from (9) that

$$1 \leq \sum_{q \in \mathcal{D}_k(N)} \sum_{0 \leq a \leq q} \frac{2C}{qN^k} \leq 4CN^{-k} |\mathcal{D}_k(N)| \ll (\log N)^{-c},$$

which is impossible when $N \rightarrow \infty$. On the other hand, one may hypothesize that the set of fractions with denominators in $\mathcal{D}_k(N)$ is distributed similarly to the set of all fractions a/q with denominators $q \leq N^k$. Under such a hypothesis, one might hope for an estimate with $|\mathcal{D}_k(N)|^{-1}$ in place of the term $(\log 3N)^c N^{-k}$ on the right side of (10) below, and such an estimate, if true, would be essentially best possible. However, upon observing that

$$|\mathcal{D}_k(N)| \geq \sum_{q_1 \leq N} \cdots \sum_{q_k \leq N} d(q_1 \cdots q_k)^{-1} \geq \left\{ \sum_{q \leq N} d(q)^{-1} \right\}^k \gg N^k (\log 3N)^{-k},$$

we will take a more cautious approach and pose the following

Question 3. Let k be a positive integer. Determine the least value of c_k such that for any real θ and any positive integer N , there exist integers $a_1, \dots, a_k, q_1, \dots, q_k$, with $1 \leq q_1, \dots, q_k \leq N$, such that

$$\left| \frac{a_1}{q_1} + \dots + \frac{a_k}{q_k} - \theta \right| \ll \frac{(\log 3N)^{c_k}}{q_1 \cdots q_k N^k}. \quad (10)$$

We leave the answer to this question to the future.

Acknowledgement. The first author would like to thank the American Institute of Mathematics for support. The second author would like to thank Jeff Vaaler for several enlightening conversations on this and related topics.

R

- [1] Tsz Ho Chan, *Finding Almost Squares II*, preprint, 2005, arXiv:math.NT/0503438.
- [2] P. Erdős, *An asymptotic inequality in the theory of numbers*, Vestnik Leningrad. Univ. **15** (1960), no. 13, 41-49 (in Russian).
- [3] G.H. Hardy and E.M. Wright, *An Introduction to the Theory of Numbers*, 5th ed., Oxford University Press, 1979.

A I M , 360 P A , P A , CA 94306, U.S.A.
E-mail address: thchan@aimath.org

D M , 1 U S , C1200, T U T
A , A , TX 78712, U.S.A.
E-mail address: kumchev@math.utexas.edu